## SYMPTOM SURVEY FORM

(Restricted to Professional Use)

TOTAL PROPERTY

PATIEN	Т _			DO	CTOR		 DATE	
INSTRUCTIONS: Number the boxes which apply to you. Use (1) for MILD symptoms (occur once or twice a year), (2) for MODERATE symptoms (occur several times a year), and (3) for SEVERE symptoms (you are aware of it almost constantly).								
					GROUP ONE		-	
1 2 3 4 5 6 7		Acid foods upset Get chilled, often "Lump" in throat Dry mouth-eyes-nose Pulse speeds after meal Keyed up — fail to calm Cuts heal slowly	8 9 10 11 12 13		Gag easily Unable to relax; startles easily Extremities cold, clammy Strong light irritates Urine amount reduced Heart pounds after retiring "Nervous" stomach	15 16 17 18 19 20	Appetite reduced Cold sweats often Fever easily raised Neuralgia-like pains Staring, blinks little Sour stomach frequent	
		1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	·		GROUP TWO		 	
21 22 23 24 25 26 27 28		Joint stiffness after arising Muscle-leg-toe cramps at night "Butterfly" stomach, cramps Eyes or nose watery Eyes blink often Eyelids swollen, puffy Indigestion soon after meals Always seems hungry; feels "lightheaded" often  Eat when nervous Excessive appetite	29 30 31 32 33 34 35 36		Digestion rapid Vomiting frequent Hoarseness frequent Breathing irregular Pulse slow; feels "irregular" Gagging reflex slow Difficulty swallowing Constipation, diarrhea alternating  GROUP THREE  Heart palpitates if meals missed or delayed	37 38 39 40 41	"Slow starter" Get "chilled" infrequently Perspire easily Circulation poor, sensitive to cold Subject to colds, asthma, bronchitis  Crave candy or coffee in afternoons	
44 45 46 47 48		Hungry between meals Irritable before meals Get "shaky" if hungry Fatigue, eating relieves "Lightheaded" if meals delayed	50 51 52		Afternoon headaches Overeating sweets upsets Awaken after few hours sleep — hard to get back to sleep	54 55	Moods of depression — "blues" or melancholy Abnormal craving for sweets or snacks	
					GROUP FOUR			
56 57		Hands and feet go to sleep easily, numbness Sigh frequently, "air	63 64		Get "drowsy" often Swollen ankles worse at night	68 69	Bruise easily, "black and blue" spots Tendency to anemia	
58		hunger" Aware of "breathing heavily"	65		Muscle cramps, worse during exercise; get "charley horses"	70 71	"Nose bleeds" frequent Noises in head, or "ringing in ears"	
59 60		High altitude discomfort Opens windows in closed room	66 67		Shortness of breath on exertion Dull pain in chest or	72	Tension under the breastbone, or feeling of "tightness",	
61 62		Susceptible to colds and fevers Afternoon "yawner"			radiating into left arm, worse on exertion		worse on exertion	

## **SYMPTOM SURVEY FORM - Page 2**

GROUP FIVE						
	73	Dizziness Dry skin Burning feet Blurred vision Itching skin and feet Excessive falling hair Frequent skin rashes Bitter, metallic taste in mouth in mornings Bowel movements painful or difficult Worrier, feels insecure	83	Feeling queasy; headache over eyes Greasy foods upset Stools light-colored Skin peels on foot soles Pain between shoulder blades Use laxatives Stools alternate from soft to watery History of gallbladder attacks or gallstones	91   92   93   94   95   96   97   97   97	Sneezing attacks Dreaming, nightmare type bad dreams Bad breath (halitosis) Milk products cause distress Sensitive to hot weather Burning or itching anus Crave sweets
				GROUP SIX		
	98	Loss of taste for meat Lower bowel gas several hours after eating Burning stomach sensations, eating relieves	101   102   103	Coated tongue Pass large amounts of foul-smelling gas Indigestion 1/2 - 1 hour after eating; may be up to 3-4 hrs.	104	Mucous colitis or "irritable bowel" Gas shortly after eating Stomach "bloating" after eating
		(A)		GROUP SEVEN		
	107   108   109   110   111   112   113   114   115   116   117   118   119   120   121	Insomnia Nervousness Can't gain weight Intolerance to heat Highly emotional Flush easily Night sweats Thin, moist skin Inward trembling Heart palpitates Increased appetite without weight gain Pulse fast at rest Eyelids and face twitch Irritable and restless Can't work under pressure	137	(C) Failing memory Low blood pressure Increased sex drive Headaches, "splitting or rending" type Decreased sugar tolerance  (D) Abnormal thirst Bloating of abdomen Weight gain around	150	(E) Dizziness Headaches Hot flashes Increased blood pressure Hair growth on face or body (female) Sugar in urine (not diabetes) Masculine tendencies (female)  (F) Weakness, dizziness Chronic fatigue
	122	(B) Increase in weight Decrease in appetite Fatigue easily Ringing in ears Sleepy during day Sensitive to cold Dry or scaly skin Constipation Mental sluggishness Hair coarse, falls out Headaches upon arising wear off during day Slow pulse, below 65 Frequency of urination Impaired hearing Reduced initiative	145	hips or waist Sex drive reduced or lacking Tendency to ulcers, colitis Increased sugar tolerance Women: menstrual disorders Young girls: lack of menstrual function	159	Low blood pressure Nails weak, ridged Tendency to hives Arthritic tendencies Perspiration increase Bowel disorders Poor circulation Swollen ankles Crave salt Brown spots or bronzing of skin Allergies - tendency to asthma Weakness after colds, influenza Exhaustion - muscular and nervous Respiratory disorders

## SYMPTOM SURVEY FORM - Page 3

FEMALE ONLY	
173 Very easily fatigued 181 Hysterectomy/ovaries 174 Premenstrual tension removed	MALE ONLY
174 Premenstrual tension removed 175 Painful menses 182 Menopausal hot 176 Depressed feelings flashes before menstruation 183 Menses scanty or 177 Menstruation excessive missed and prolonged 184 Acne, worse at 178 Painful breasts menses 179 Menstruate too frequently 185 Depression of 180 Vaginal discharge long standing	186 Prostate trouble  187 Urination difficult or dribbling  188 Night urination frequent  189 Depression
IMPORTANT  TO THE PATIENT: Please list below the five main physical complaints you have in order of their importance:  1	190 Pain on inside of legs or heels  191 Feeling of incomplete bowel evacuation  192 Lack of energy  193 Migrating aches and pains  194 Tire too easily  195 Avoids activity  196 Leg nervousness at night  197 Diminished sex drive