



# ARRAY

## 5

## MULTIPLE AUTOIMMUNE REACTIVITY SCREEN™



2 mL Serum

### Antigens Tested (IgG + IgA Combined)

- Parietal Cell + ATPase
- Intrinsic Factor
- ASCA + ANCA
- Tropomyosin
- Thyroglobulin
- Thyroid Peroxidase (TPO)
- 21 Hydroxylase (Adrenal Cortex)
- Myocardial Peptide
- Alpha-Myosin
- Phospholipid
- Platelet Glycoprotein
- Ovary/Testis\*
- Fibulin
- Collagen Complex
- Arthritic Peptide
- Osteocyte
- Cytochrome P450 (Hepatocyte)
- Insulin + Islet Cell Antigen
- Glutamic Acid Decarboxylase 65 (GAD 65)
- Myelin Basic Protein
- Asialoganglioside
- Alpha + Beta Tubulin
- Cerebellar
- Synapsin

\*Ovary and Testis are tested together to avoid any confusion arising out of potential cross-reactivity.

### CLINICAL USE:

- Measure Predictive Antibodies, some of which can appear up to ten years before the clinical onset of disease.
- Economically and efficiently assess possible tissue damage to multiple organs of the body.

### RECOMMENDED FOR PATIENTS WHO:

- Have a diagnosed autoimmune disorder, as they commonly develop additional autoimmune conditions.
- Have chronic increased intestinal permeability, which is the gateway for environmentally-induced autoimmune disorders.

 **CYREX**<sup>®</sup>  
Laboratories  
[www.JoinCyrex.com](http://www.JoinCyrex.com)

# MULTIPLE AUTOIMMUNE REACTIVITY SCREEN™

